

**Supplementary Table 1.** Awareness of suicide according to the severity of depressive symptoms

Variables	No depression (N=292)	Mild depression (N=196)	Moderate depression (N=127)	Severe depression (N=22)
Importance of suicide prevention				
Agree	260 (89.0)	156 (79.6)	84 (66.1)	12 (54.5)
Unsure	21 (7.2)	21 (10.7)	30 (23.6)	6 (27.3)
Indifferent	4 (1.4)	14 (7.1)	11 (8.7)	1 (4.5)
Disagree	7 (2.4)	5 (2.6)	2 (1.6)	3 (13.6)
Possibility of suicide prevention				
Absolutely impossible	7 (2.4)	2 (1.0)	6 (4.7)	2 (9.1)
Impossible	8 (2.7)	21 (10.7)	24 (18.9)	6 (27.3)
Unsure	79 (27.1)	77 (39.3)	44 (34.6)	6 (27.3)
Possible	122 (41.8)	77 (39.3)	49 (38.6)	6 (27.3)
Absolutely possible	76 (26.0)	19 (9.7)	4 (3.1)	2 (9.1)
Frequency of contacting someone				
Almost everyday	70 (24.0)	32 (16.3)	17 (13.4)	0 (0.0)
Once a week	103 (35.3)	59 (30.1)	46 (36.2)	4 (18.2)
1–2 times a month	86 (29.5)	55 (28.1)	36 (28.3)	6 (27.3)
1–2 times in 6 months	24 (8.2)	28 (14.3)	15 (11.8)	1 (4.5)
1–2 times a year	4 (1.4)	10 (5.1)	2 (1.6)	3 (13.6)
Almost none	5 (1.7)	12 (6.1)	11 (8.7)	8 (36.4)
SSI sum	2.76±2.87	5.36±4.47	11.49±6.76	16.95±7.67

Values are presented as N (%) or mean±standard deviation. SSI, Scale for suicidal ideation