

Supplementary Table 2. Differences of MMPI-2 supplementary scales between stimulants and cannabis users

MMPI-2 supplementary scales	Stimulants (N=18)	Cannabis (N=60)	F*
A	49.0±14.7	43.9±11.9	0.84
R	51.2±9.6	49.7±8.9	0.23
Es	51.8±12.1	55.8±10.6	0.07
Do	49.2±11.6	52.0±9.5	0.53
Re	48.3±14.2	51.0±11.5	0.03
Mt	50.8±16.4	44.0±13.7	1.25
PK	53.3±16.7	45.3±13.8	1.31
MDS	51.1±14.8	45.7±11.2	0.79
Ho	46.8±8.3	45.8±12.2	0.05
O-H	55.6±6.6	51.4±12.0	1.17
MAC-R	55.3±10.9	54.4±10.3	0.17
AAS	51.4±14.1	49.3±11.7	0.83
APS	55.2±13.5	53.2±9.9	0.12
GM	50.7±14.6	59.3±9.6	0.83
GF	51.9±9.6	48.6±8.9	0.32

Values are presented as mean±standard deviation. *covariates: age, sex, marital status, and past psychiatry history. A, Anxiety; R, Repression; Es, Ego-Strength; Do, Dominance; Re, Social Responsibility; Mt, College Maladjustment; PK, Post-traumatic Stress disorder; MDS, Marital Distress Scale; Ho, Hostility; O-H, Overcontrolled Hostility; MAC-R, MacAndrew Alcoholism Scale; AAS, Addiction Acknowledgment Scale; APS, Addiction Potential Scale; GM, Gender role-Masculine; GF, Gender role-Feminine