

**Supplementary Table 5.** Pearson's correlation analyses among continuous variables in patients with panic disorder

Continuous variables	1	2	3	4	5	6	7	8	9	10	11	12	13	14
Age (1)	-													
ETISR-SF														
Total sum of all subtypes (2)	-0.202***	-												
General (3)	-0.072	0.642***	-											
Physical (4)	-0.150**	0.644***	0.296***	-										
Emotional (5)	-0.239***	0.712***	0.348***	0.372***	-									
Sexual (6)	-0.020	0.478***	0.294***	0.160**	0.274***	-								
Coping strategies														
Problem-focused (7)	0.112*	-0.003	0.060	0.012	-0.035	0.065	-							
Emotion-focused (8)	-0.160**	0.219***	0.146**	0.204***	0.267***	0.135**	0.393***	-						
APPQ														
Total sum of all subtypes (9)	-0.124*	0.204***	0.143**	0.055	0.310***	0.100*	-0.007	0.322***	-					
Agoraphobia (10)	-0.120*	0.138***	0.107*	-0.008	0.261***	0.061	0.032	0.277***	0.911***	-				
Social phobia (11)	-0.163**	0.250***	0.154**	0.136**	0.320***	0.136**	-0.088	0.337***	0.860***	0.670***	-			
Interoceptive fear (12)	-0.004	0.104**	0.073	-0.007	0.197***	0.069	0.048	0.198***	0.841***	0.697***	0.553***	-		
PDSS total scores (13)	-0.235***	0.112*	0.067	0.090	0.153**	0.003	-0.007	0.259***	0.460***	0.456***	0.340***	0.355***	-	
SSI total scores (14)	-0.224***	0.270***	0.139**	0.201***	0.350***	0.140**	-0.066	0.372***	0.446***	0.385***	0.454***	0.290***	0.346***	-

\*p<0.05; \*\*p<0.01; \*\*\*p<0.001. ETISR-SF, Early Trauma Inventory Self-Report-Short Form; APPQ, Albany Panic and Phobia Questionnaire; PDSS, Panic Disorder Symptom Severity; SSI, Scale for Suicidal Ideation