Supplementary Table 3. Differences between missing and complete cases at the 6-month assessment

Variables	Missing cases (N=91)	Complete cases (N=63)	t/χ^2	p
Sociodemographic				
Age (yr)	50.57±10.51	51.10±10.24	-0.307	0.759
Marital status			1.106	0.575
Currently married or cohabitating	68 (74.7)	43 (68.3)		
Never married	13 (14.3)	13 (20.6)		
Separated, divorced, or widowed	10 (11.0)	7 (11.1)		
Living status			0.786	0.375
With family or nonfamily	81 (89.0)	53 (84.1)		
Alone	10 (11.0)	10 (15.9)		
Education			1.562	0.211
College or higher	61 (67.0)	36 (57.1)		
High school or lower	30 (33.0)	27 (42.9)		
Employment			0.326	0.568
Employed	32 (35.2)	25 (39.7)		
Unemployed	59 (64.8)	38 (60.3)		
Cancer-related				
Time passed since index cancer diagnosis (m)	43.01±57.22	40.63±43.57	0.144	0.781
Surgery			3.351	0.067
No	65 (71.4)	53 (84.1)		
Yes	26 (28.6)	10 (15.9)		
Chemotherapy			1.448	0.229
No	66 (72.5)	51 (81.0)		
Yes	25 (27.5)	12 (19.0)		
Radiotherapy			-	>0.999
No	84 (92.3)	59 (93.7)		
Yes	7 (7.7)	4 (6.3)		
Other therapy			1.131	0.287
No	44 (48.4)	25 (39.7)		
Yes	47 (51.6)	38 (60.3)		
Psychiatric-related				
Psychiatric disorder			-	0.145
No	4 (4.4)	0 (0.0)		
Yes	87 (95.6)	63 (100.0)		
History of psychiatric admission			-	>0.999
No	85 (93.4)	59 (93.7)		
Yes	6 (6.6)	4 (6.3)		
History of suicide attempt			0.135	0.713
No	73 (80.2)	49 (77.8)		
Yes	18 (19.8)	14 (22.2)		
Familial history of psychiatric disorder			2.388	0.122
No	78 (85.7)	59 (93.7)		
Yes	13 (14.3)	4 (6.3)		
Grit				
Consistency of interests	20.48±4.45	19.86±4.62	0.845	0.399
Perseverance of effort	19.10±4.38	18.92±4.29	0.250	0.803
Psychiatric				
PHQ-9*	9.14±6.26	9.41±6.10	-0.266	0.791
GAD-7†	6.37±5.24	7.79±5.56	-1.613	0.109
Pain				
PPI	1.52±1.04	1.43±0.911	0.543	0.588

Data are presented as mean±standard deviation or number (%). *the number of missing cases is 19; †the number of missing cases is 20. PHQ-9, Patient Health Questionnaire-9; GAD-7, Generalized Anxiety Disorder-7; PPI, Present Pain Index.